

Breaking the cycle: Reforming pesticide regulation to protect pollinators

Adrian II Fisher, Rafaela Tadei, May Berenbaum, James Nieh, Harry Siviter, James Crall , Jordan R. Glass , Felicity Muth , Ling-Hsiu Liao, Kirsten Traynor, Nicole DesJardins, Roberta Nocelli , Noa Simon-Delso and Jon F. Harrison 

Adrian Fisher II (afishe16@asu.edu), Jordan R. Glass (jrglass@asu.edu), Nicole DesJardins (ndesjard@asu.edu), and Jon F. Harrison (j.harrison@asu.edu) are affiliated with the School of Life Sciences at Arizona State University, in Tempe, Arizona, in the United States. Rafaela Tadei (rafaelatadei@gmail.com) is affiliated with the São Paulo State University, in Rio Claro, Brazil. May Berenbaum (maybe@illinois.edu) is affiliated with the University of Illinois at Urbana-Champaign, in Urbana, Illinois, in the United States. James Nieh (jnieh@ucsd.edu) is affiliated with the University of California, in San Diego, California, in the United States. Harry Siviter (harry.siviter@austin.utexas.edu) is affiliated with the University of Texas at Austin, in Austin, Texas, in the United States, and with the University of Bristol, in Bristol, England, in the United Kingdom. James Crall (james.crall@wisc.edu) is affiliated with the University of Wisconsin-Madison, in Madison, Wisconsin, in the United States. Felicity Muth (felicity.muth@austin.utexas.edu) is affiliated with the University of Texas at Austin, in Austin, Texas, in the United States. Ling-Hsiu Liao (liao19@illinois.edu) is affiliated with the University of Illinois at Urbana-Champaign, in Urbana, Illinois, in the United States. Kirsten Traynor (kirsten.traynor@uni-hohenheim.de) is affiliated with the University of Hohenheim, in Stuttgart, Germany. Roberta Nocelli (roberta@ufscar.br) is affiliated with the Federal University of São Carlos, in Araras, Brazil. Noa Simon-Delso (simon@bee-life.eu) is affiliated with BeeLife European Beekeeping Coordination, in Louvain la Neuve, Belgium.

Abstract

Over decades, pesticide regulations have cycled between approval and implementation, followed by the discovery of negative effects on nontarget organisms that result in new regulations, pesticides, and harmful effects. This relentless pattern undermines the capacity to protect the environment from pesticide hazards and frustrates end users that need pest management tools. Wild pollinating insects are in decline, and managed pollinators such as honey bees are experiencing excessive losses, which threatens sustainable food security and ecosystem function. An increasing number of studies demonstrate the negative effects of field-realistic exposure to pesticides on pollinator health and fitness, which contribute to pollinator declines. Current pesticide approval processes, although they are superior to past practices, clearly continue to fail to protect pollinator health. In the present article, we provide a conceptual framework to reform cyclical pesticide approval processes and better protect pollinators.

Keywords: pollinators, pesticides, policy, regulation, sustainability

Trevan (1927) proposed the median lethal dose (LD₅₀) as one useful measure for standardizing drug dosages, but in the intervening 95 years, it has become the foundation for assessing the acute toxicity and environmental safety of agrochemicals, including pesticides. But there is almost a century of evidence that the LD₅₀ metric has systematically failed to identify critical negative impacts (just short of lethality) on beneficial insects that provide important ecosystem services—most notably, pollination. Reliance on lethality assessments has created a repeating cycle in which pesticides are approved and used until detrimental impacts on nontarget species are discovered, with ensuing restrictions, followed by the development of new replacement pesticides that are often subsequently discovered to also be hazardous (figure 1). Many studies have demonstrated that numerous approved pesticides have significant negative sublethal effects on the behavior, physiology, and reproduction of managed and wild pollinators, both of which are economic contributors to agriculture, at field-realistic levels (Desnaux et al. 2007, Siviter et al. 2021a, Tosi et al. 2022). In addition, because biological processes tend to be conserved within a taxon, the documentation of harmful sublethal effects of approved pesticides on managed or model pollinators almost certainly means that the pesticides are also dangerous for nonmodel, nonmanaged pollinators (Woodcock et al. 2016).

Many wild pollinator species are experiencing population declines, whereas managed pollinators, most notably honey bees

(*Apis mellifera*), are experiencing economically unsustainable regional losses, threatening sustainable food security, ecosystem function, and human health (Smith et al. 2015, Reilly et al. 2020). Although the relative importance of pesticides to insect decline remains unclear, there is unequivocal evidence that pollinator safety across many landscapes is put at risk by current pesticide use practices (Siviter et al. 2021b). Long-term preservation of both biodiversity and agricultural sustainability demands a major revision of pesticide approval systems (Drivdal and van der Sluijs 2021). In the present article, we suggest a pathway to develop pesticide approval systems to anticipate and prevent inadvertent adverse impacts of pesticides on nontarget species. We first highlight the limitations of current pesticide regulation protocols and then suggest steps for improving pesticide regulation, including harnessing recent technological advances that overcome the limitations of the current regulation system.

Current regulatory approaches and inadequacies

Pesticide toxicity is currently tested with a tiered approach that is intended to allow approval of only those chemicals at low risk of causing adverse impacts on the biotic environment (EPA 2014a, EFSA et al. 2023). The risks presented by pesticides to pollinators that encounter them have long been assessed using honey bees as

Received: April 24, 2023. Revised: August 29, 2023. Accepted: September 8, 2023

© The Author(s) 2023. Published by Oxford University Press on behalf of the American Institute of Biological Sciences. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. Comparison of current pesticide regulatory practices and a suggested model for improvement. (a) The current pesticide assessment process includes approval of a pesticide on the basis of low mortality of larvae and caged honey bee adults in contact and oral exposure tests. The current practices allow bypassing of spraying in outdoor cages and field tests (no cages) instead using only first tier tests to indicate low risk on the basis of short-term LD₅₀ evaluations. In addition, current practices in most countries do not include postapproval monitoring that could result in the banning of a pesticide because of its nontarget effects. (b) The suggested improved pesticide approval and monitoring process includes requirements for multiple levels of testing with field- and taxon-relevant exposure conditions and risk assessment before a pesticide is approved. This process would also require assessing pesticide toxicity for diverse pollinator species, testing of sublethal effects, and synergistic interactions between pesticides. In addition, our proposed model includes post-approval monitoring allowing for enhanced efficiency in detecting unanticipated negative effects.

a proxy for the thousands of insect species that contribute to plant pollination. In the United States and Brazil, exposures are estimated with the Bee-REX model, which incorporates worst-case scenario exposure events (EPA 2014b, Cham et al. 2020). In Europe, various exposure models have been proposed depending on the pesticide application way (i.e., contact and dietary models), leading to various predictive exposure concentrations that result from the consideration of parameters such as the application rate, exposure factors, the body surface factor, or the food consumption (EFSA et al. 2023). These are then compared with LD₅₀ values under laboratory conditions. Many pesticides undergo only a first-tier screen with honey bee adults and larvae under laboratory conditions. Next-tier studies can include quantification of pesticide residues in pollen and nectar collected by bees and pesticide effects on physiology and behavior of lab-reared bees. These next-tier studies are conducted only if the calculated risk quotients from the laboratory studies exceed the levels of concern or if other data indicate behavioral or fitness consequences of the pesticide during the approval process (CFR40C § 158.630; EPA 1996). Similarly, the highest tier studies of effects of pesticides on pollinators in the field are conducted only if concerns arise from lower tier studies (European Commission 2011, EPA 2014a, Sanchez-Bayo and Goka 2014). The effects of pesticide toxicity have become more apparent over time because of development of methodologies focusing on more subtle observations than mortality. This suggests that current exposure duration standards for tier 1 tests do not allow for assessments of chronic nor sublethal effects to clearly lay out proper boundaries for protecting pollinators (Sanchez-Bayo and Tennekes 2020).

Unfortunately, this system has repeatedly failed to identify sublethal injurious effects on pollinators. Field-realistic concentrations of numerous approved fungicides, herbicides, and insecticides demonstrably compromise pollinator health (Iwasaki

and Hogendoorn 2021). At the individual level, exposure to certain pesticides can undermine learning and navigation abilities, digestive functioning, brood care, flight performance, and longevity (Tosi et al. 2017, Siviter et al. 2018, Kenna et al. 2019, Fisher et al. 2021, Fischer et al. 2023). At the colony level, sublethal consequences of field-realistic pesticide exposures include disruption of thermoregulatory capacity, reduction in colony size, suppressed male and female reproduction, increased brood disease, and reduced overwintering survival (Whitehorn et al. 2012, Crall et al. 2018, Traynor et al. 2021). Although the majority of such studies have been performed with honey bees as the model nontarget species, bumble bees, stingless bees, and solitary bees can experience comparable or even more severe adverse impacts (Artz and Pitts-Singer 2015, Crall et al. 2018).

Failures to identify and document the adverse ecological impacts of pesticides arise in multiple ways. One clear type of failure has been the reliance on laboratory lethality (i.e., LD₅₀) assays for short exposures (European Commission 2013). Short-term LD₅₀s are problematic because pollinators generally experience exposure to agrochemicals for much longer than a few days, and the toxicities of agrochemicals to pollinators have been well-demonstrated to increase with duration of exposure (Simon-Delso et al. 2018, Sánchez-Bayo and Tennekes 2020). Also, LD₅₀s characterize a dose that is highly toxic, killing 50% of the population, and the relationships between such a dose and the dose that will have a minimal effect on pollinators are generally poorly characterized. Current approval procedures do not as a matter of course include assessments of sublethal effects or field-realistic outcomes (European Commission 2002, EPA 2014a). Characterizing the links between sublethal effects on individual performance (e.g., learning and navigation abilities, digestive functioning, and thermoregulation), pollinator fitness typically requires long assessment periods under field-realistic exposure regimes. For eusocial pollinators,

such as honey bees and bumble bees, toxicity tests on individual workers, most of which are sterile (nonreproductive), do not predict colony-level effects mediated by the responses of the few reproductive individuals in the colony to pesticide exposure or by colony-level feedback regulation. Another limitation of current procedures is the narrow focus on isolated active ingredients, whereas, in reality, many pesticide formulations contain mixtures of chemicals that can interact with other factors or contain “inert ingredients” that are, in fact, toxic (Straw et al. 2022). Moreover, pollinators may be exposed simultaneously to diverse types of pesticides, as well as to secondary stressors (e.g., disease, poor nutrition) that may sometimes have strong interactive or additive effects with pesticides in the field.

The near-exclusive reliance on honey bees as a model species in toxicity testing is problematic, as has been demonstrated by multiple direct sensitivity comparisons across bee species (EFSA Panel on Plant Protection Products and their Residues 2012, Arena and Sgolastra 2014, Gradish et al. 2019). Pollinators differ widely in life history and physiology, resulting in tremendous variation in exposure routes and susceptibilities (Hladik et al. 2016, Boyle et al. 2019, Gradish et al. 2019). Finally, challenges in estimating the persistence of pesticides in the environment have caused insufficient estimations of exposure (Traynor et al. 2021). As a prime example, exposure to wind-blown dust from seeds coated with neonicotinoids was not predicted to be a major route of exposure when these pesticides were approved (Sgolastra et al. 2020, Greatti et al. 2003).

Solutions

Over the past century of pesticide testing, improvements have been made in better describing the routes of pesticide exposure and developing methodologies to reveal their effects. That said, maintaining such a slow pace of advancement is insufficient for the task of protecting pollinators. We offer a five-part solution:

Thorough laboratory screening

The current reliance on mortality levels (i.e., LD₅₀) within a short time frame (48 hours) as the standard for assessing pesticide toxicity to insect pollinators is the first limitation in adjusting procedural inadequacies (OECD 1998). For other model organisms, LD₅₀ testing has been discouraged and replaced with other methods that monitor lethality and various outcomes besides mortality for animal subjects (Erhirhie et al. 2018). Recently, the European Food Safety Authority (EFSA) reconsidered the hazard parameters in experimental studies and proposed to extract more information from the toxicological tests, considering the whole dose–response curves (from which other toxicological endpoints could be extrapolated—no observed effect concentration or LD_x—calculating the existence of time-reinforced toxicity or integrating sublethal effects and mixture toxicity; EFSA et al. 2023). Such an approach with pollinators would increase test sensitivity and aid in addressing the disconnection between laboratory and field results where outcomes of exposure include a variety of sublethal effects at low concentrations. In addition, the current limits for oral toxicity assessments range from 48 hours to 10 days, which does not reflect field-relevant exposure or allow for adequate evaluation of resulting outcomes (OECD 1998, OECD 2017, Simon-Delso et al. 2018). Assessments of mortality should discard rigid timeframes, instead allowing the lifespan and natural history of test organisms or the persistence of pesticide residues in nature to factor into expo-

sure and overall test duration (Tosi et al. 2021). In general, toxicological testing for risk assessment purposes should be prolonged in time. Furthermore, decision-makers should consider the methodological developments proposed by the EFSA (EFSA et al. 2022, 2023), which may allow for progress in describing the pesticide toxicological profile and its risks to pollinators. On the basis of this information, decision-makers will be able to base their approval decisions on more relevant information.

Relevant sublethal testing

The approval process must include assessments of sublethal effects on individuals and colony-level effects such as reproduction under a range of possible concentrations and realistic exposure times for pollinators. Adjusted toxicity standards using LD₁₀ values or NOEL (no observed effect levels) would allow for reduced lethality and more careful discernment of nonlethal detrimental effects of exposure to establish protective standards. Studies that include measurement of pesticide concentrations in resources collected and used by pollinators are needed for calculating field-realistic exposures (Linguadoca et al. 2021). For eusocial pollinators, colony-level testing must be mandatory to determine safe-use levels because effects on colony reproduction or colony-level behavioral interactions cannot be assessed fully by testing individuals in the laboratory. In addition, prolonged laboratory tests using field-realistic concentration and exposure time can be useful to show physiological, behavioral, and other sublethal effects as a proxy for determining the effects on pollinator population growth and development (Simon-Delso et al. 2018). Also, tests targeted at uncovering underlying mechanistic effects of exposure can inform physiological outcomes and behavioral observations. Methods for measuring sublethal or colony-level effects that are not prohibitively expensive are available and well-established for many taxa. Technological advances are making scalable, cost-effective studies of the sublethal and synergistic impacts of agrochemicals feasible. In particular, the rapid emergence of computer vision tools for behavior provides new opportunities for developing reliable, standardized, and economical screening protocols (Crall et al. 2018, Høye et al. 2020).

Testing diverse species

In view of the physiological and ecological diversity that exists across pollinator taxa, testing procedures must include regionally appropriate managed and native bees and other pollinator species likely to be exposed to the pesticide. In many tropical regions, stingless bees are primary pollinators, and they have been shown to respond differently to multiple pesticides than honey bees (Cham et al. 2018). Similarly, multiple native bee species have different life histories than honey bees, including soil nesting and varying levels of sociality and, therefore, experience differing exposure routes and risks (Boyle et al. 2019, Franklin and Raine 2019, Sgolastra et al. 2019). The approval process must assess the extent to which native pollinators are likely to experience pesticide exposure, so that toxicity levels can be measured with field-realistic exposures. In the revised guidance to pollinator risk assessment from the EFSA, at least one species of bumble bee and one species of solitary bee have been proposed for inclusion in the pesticide regulation (EFSA et al. 2023). However, a number of other insect pollinators with differing biology and vulnerability to pesticides may exist, which should, ideally, be considered in the future improvement of the pesticide risk assessment (EFSA et al. 2022). In addition, nonbee insect pollinators provide substantial contributions to pollination, can experience similar agrochemical

exposure, and should be considered for toxicity testing (Rader et al. 2015, Uhl and Brühl 2019).

Mandatory field-realistic testing

The regulatory community should implement and mandate thorough and cost-effective safety testing mechanisms to ensure that approved pesticides are safe for pollinators under realistic environmental conditions. Although they are few in number, several field tests of pesticide exposure demonstrate numerous negative outcomes for multiple pollinator species (Rundlöff et al. 2015, Tsvetkov et al. 2017, Woodcock et al. 2017, Olaya-Arenas et al. 2020, Fisher et al. 2022). The repeated failures of the current system for predicting nontarget pesticide toxicity demonstrate that laboratory determinations of LD₅₀ values are insufficient. Conceivably, improvements in lower-tier testing as described above might ameliorate the need for field testing. However, given that environmental conditions usually differ dramatically from lab conditions, that environmental conditions such as temperature affect toxicity (Kenna et al. 2023), and that for eusocial insects, colony-level testing is critical to assess effects on reproduction, mandating higher-tier field-realistic testing before approval decisions are made could significantly lower threats to pollinator safety (figure 1). In addition, pesticides should be tested as they are formulated and sold for real-world use to account for the potential effects of inert ingredients and the synergisms between formulation components. Field tests can improve the interpretation of results in combination with the lethal and sublethal tests proposed in the present article, their results being considered as complementary to the laboratory tests rather than as probatory of risk or no risk (figure 1b)

Postapproval monitoring (fitopharmacovigilance) and reassessment

It is difficult for scientists to anticipate all important field fates of pesticides and their interactions, so postapproval monitoring and reassessment of provisionally approved pesticides, as is routinely performed for pharmaceuticals (Milner and Boyd 2017), must be implemented. This process should include surveys of both managed and unmanaged pollinators exposed to pesticides applied in the approved manner (figure 1).

Developing an effective system for testing pesticide safety will require the engagement of many more stakeholders, including those with expertise in chemical environmental degradation and pesticide residues in realistic environmental conditions, as well as those with expertise in pollinator behavior, physiology, and natural history. A layered approach with multiple testing levels accounting for underlying mechanisms paired with clear measures of behavior and fitness will provide policymakers with the best data for making pesticide approval decisions (Ankley et al. 2010, Sponsler et al. 2019). Overhauling the current pesticide approval system will require significant political and economic resolve. Recently, the United States Geological Survey (USGS) unveiled plans to reduce the extent of their pesticide monitoring and frequency of pesticide use reporting as part of the Pesticide National Synthesis Project (USGS 2023). Such measures follow prior reductions in the last decade that also scaled back the comprehensiveness of the USGS's pesticide monitoring program (Hitaj et al. 2020, Douglas 2023). Although it was once a valuable tool for researchers, this reconfiguration of the USGS pesticide database constitutes a major setback for investigations into pesticide use and pollinator declines (Gewin 2023). Legislative input in maintaining pesticide monitoring databases and implementing stringent postapproval

pesticide evaluations is therefore essential for enhanced pollinator protection. The consequences of reduced pesticide monitoring and inaction in changing toxicity testing could have wide spread ramifications, especially if the losses of managed or wild pollinators increase. Pollinator insufficiency could compromise food security for vulnerable populations around the world (Steffan-Dewenter et al. 2005).

Acknowledgments

We thank the Entomological Society of America for providing the platform on which the authors initially met and discussed common themes of pesticide effects on pollinators.

References cited

- Ankley GT, et al. 2010. Adverse outcome pathways: A conceptual framework to support ecotoxicology research and risk assessment. *Environmental Toxicology and Chemistry* 29: 730–741.
- Arena M, Sgolastra F. 2014. A meta-analysis comparing the sensitivity of bees to pesticides. *Ecotoxicology* 23: 324–334.
- Artz DR, Pitts-Singer TL. 2015. Effects of fungicide and adjuvant sprays on nesting behavior in two managed solitary bees, *Osmia lignaria* and *Megachile rotundata*. *PLOS ONE* 10: e0135688.
- Boyle NK, et al. 2019. Workshop on pesticide exposure assessment paradigm for non-Apis bees: Foundation and summaries. *Environmental Entomology* 48: 4–11.
- Cham KO, et al. 2018. Pesticide exposure assessment paradigm for stingless bees. *Environmental Entomology* 48: 36–48.
- Cham KO, Rebelo RM, Oliveira RP, Ferro AA, Silva FECV, Borges LO, Saretto COSD, Tonelli CAM, Macedo TC. 2020. Avaliação de Risco Ambiental de Agrotóxicos para Abelhas IBAMA/Diqua.
- Crall JD, et al. 2018. Neonicotinoid exposure disrupts bumblebee nest behavior, social networks, and thermoregulation. *Science* 362: 683–686.
- Desnoux N, Decourtye A, Delpuech JM. 2007. The sublethal effects of pesticides on beneficial arthropods. *Annual Review of Entomology* 52: 81–106.
- Douglas M. 2023. An essential U.S. pesticide database is disappearing. *Entomology Today* (30 June 2023). <https://entomologytoday.org/2023/06/29/pesticide-national-synthesis-project-essential-database-disappearing>.
- Drivdal L, van der Sluijs JP. 2021. Pollinator conservation requires a stronger and broader application of the precautionary principle. *Current Opinion in Insect Science* 46: 95–105.
- [EFSA] European Food Safety Authority et al. 2022. Theme (concept) paper: Advancing the environmental risk assessment of chemicals to better protect insect pollinators (IPol-ERA). *EFSA Journal* 19: E200505E.
- [EFSA] European Food Safety Authority et al. 2023. Revised guidance on the risk assessment of plant protection products on bees (*Apis mellifera*, *Bombus* spp. and solitary bees). *EFSA Journal* 21: e07989.
- EFSA Panel on Plant Protection Products and their Residues. 2012. Scientific opinion on the science behind the development of a risk assessment of plant protection products on bees (*Apis mellifera*, *bombus* spp., and solitary bees). *EFSA Journal* 10: 2668.
- [EPA] Environmental Protection Agency. 1996. *Ecological Effects Test Guidelines OPPTS 850.3040: Field Testing for Pollinator*. EPA.
- [EPA] Environmental Protection Agency. 2014a. *Guidance for Assessing Pesticide Risks to Bees*. EPA.
- [EPA] Environmental Protection Agency. 2014b. *Guidance for Assessing Pesticide Risks to Bees*. EPA.

- Erhirhie EO, Ihekwereme CP, Ildigwe EE 2018. Advances in acute toxicity testing: Strengths, weaknesses and regulatory acceptance. *Interdisciplinary Toxicology* 11: 5–12.
- European Commission. 2002. Guidance document on Terrestrial Ecotoxicology under Council Directive 91/414/EEC. European Commission. Report no. Sanco/10329/2002 rev 2.
- European Commission. 2011. Commission Regulation (EU) No 546/2011 of 10 June 2011 implementing Regulation (EC) No 1107/2009 of the European Parliament and of the Council as regards uniform principles for evaluation and authorisation of plant protection products text with EEA relevance. *Official Journal of the European Union* 155: 127–175.
- European Commission. 2013. Commission Regulation (EU) No 283/2013 of 1 March 2013 setting out the Data Requirements for active substances, in accordance with Regulation (EC) No 1107/2009 of the European Parliament and of the Council Concerning the Placing of Plant Protection Products on the market. *Official Journal of the European Union* 93: 1–84.
- Fisher A II, DeGrandi-Hoffman G, Smith BH, Johnson M, Kaftanoglu O, Cogely T, Fewell JH, Harrison JF. 2021. Colony field test reveals dramatically higher toxicity of a widely-used mitotic fungicide on honey bees (*Apis mellifera*). *Environmental Pollution* 269: 115964.
- Fisher A II, Glass JR, Ozturk C, Desjardins N, Raka Y, DeGrandi-Hoffman G, Smith BH, Fewell JH, Harrison JF. 2022. Seasonal variability in physiology and behavior affect the impact of fungicide exposure on honey bee (*Apis mellifera*) health. *Environmental Pollution* 311: 120010.
- Fischer LR, Ramesh D, Weidenmüller A. 2023. Sub-lethal but potentially devastating: The novel insecticide flupyradifurone impairs collective brood care in bumblebees. *Science of the Total Environment* 903: 166097.
- Franklin EL, Raine NE. 2019. Moving beyond honeybee-centric pesticide risk assessments to protect all pollinators. *Nature Ecology and Evolution* 3: 1373–1375.
- Gewin V. 2023. Move to change how U.S. tracks pesticide use sparks protest. *Science* (30 May 2023). www.science.org/content/article/move-change-how-u-s-tracks-pesticide-use-sparks-protests.
- Gradish AE, et al. 2019. Comparison of pesticide exposure in honey bees (Hymenoptera: Apidae) and bumble bees (Hymenoptera: Apidae): Implications for risk assessments. *Environmental Entomology* 48: 12–21.
- Greatti M, Sabatini AG, Barbattini R, Rossi S, Stravisi A 2003. Risk of environmental contamination by the active ingredient imidacloprid used for corn seed dressing. Preliminary results. *Bulletin of Insectology* 56: 69–72.
- Hitaj C, Smith DJ, Code A, Wechsler S, Esker PD, Douglas MR 2020. Sowing uncertainty: What we do and don't know about the planting of pesticide-treated seed. *BioScience* 70: 390–403.
- Hladik ML, Vandever M, Smalling KL 2016. Exposure of native bees foraging in an agricultural landscape to current-use pesticides. *Science of the Total Environment* 542: 469–477.
- Høyve TT, Årje J, Bjerke K, Hansen OLP, Iosifidis A, Leese F, Mann HMR, Meissner K, Melvad C, Raitoharju J 2020. Deep learning and computer vision will transform entomology. *Proceedings of the National Academy of Sciences* 118: e2002545117.
- Iwasaki JM, Hogendoorn K 2021. Non-insecticide pesticide impacts on bees: A review of methods and reported outcomes. *Agriculture, Ecosystems, and Environment* 314: 107423.
- Kenna D, Cooley H, Pretelli I, Rodrigues AR, Gill SD, Gill RJ 2019. Pesticide exposure affects flight dynamics and reduces flight endurance in bumblebees. *Ecology and Evolution* 9: 5637–5650.
- Kenna D, Graystock P, Gill RJ 2023. Toxic temperatures: Bee behaviours exhibit divergent pesticide toxicity relationships with warming. *Global Change Biology* 29: 2981–2998.
- Linguadoca A, Rizzi C, Villa S, Brown MJF. 2021. Sulfoxaflor and nutritional deficiency synergistically reduce survival and fecundity in bumblebees. *Science of the Total Environment* 795: 148680.
- Main AR, Hladik ML, Webb EB, Goyne KW, Mengel D 2020. Beyond neonicotinoids: Wild pollinators are exposed to a range of pesticides while foraging in agroecosystems. *Science of the Total Environment* 742: 140436.
- Milner AM, Boyd IL. 2017. Toward pesticidovigilance. *Science* 357: 1232–1234.
- [OECD] Organisation for Economic Cooperation and Development. 1998. OECD Guidelines for the Testing of Chemicals, section 2: Test no. 213: Honeybees, Acute Oral Toxicity Test. OECD. <https://doi.org/10.1787/9789264070165-en>.
- [OECD] Organisation for Economic Cooperation and Development. 2017. OECD Guidelines for the Testing of Chemicals, section 2: Test no. 245: Honey Bee (*Apis mellifera* L.), Chronic Oral Toxicity Test (10-Day Feeding). OECD. <https://doi.org/10.1787/9789264284081-en>.
- Olaya-Arenas P, Scharf ME, Kaplan I. 2020. Do pollinators prefer pesticide-free plants? An experimental test with monarchs and milkweeds. *Journal of Applied Ecology* 57: 2019–2030.
- Rader R, Bartomeus I, Garibaldi LA, Woyciechowski M. 2015. Non-bee insects are important contributors to global crop pollination. *Proceedings of the National Academy of Sciences* 113: 146–151.
- Reilly JR, et al. 2020. Crop production in the USA is frequently limited by a lack of pollinators. *Proceedings of the Royal Society B* 287: 20200922.
- Rundlöff M, et al. 2015. Seed coating with a neonicotinoid insecticide negatively affects wild bees. *Nature* 521: 77–80.
- Sánchez-Bayo F, Goka K. 2014. Pesticide residues and bees: A risk assessment. *PLOS ONE* 9: e94482.
- Sánchez-Bayo F, Tennekes HA. 2020. Time-cumulative toxicity of neonicotinoids: Experimental evidence and implications for environmental risk assessments. *International Journal of Environmental Research and Public Health* 17: 1629.
- Sgolastra F, Hinarejos S, Pitts-Singer TL, Boyle NK, Joseph T, Lückmann J, Raine NE, Singh R, Williams NM, Bosch J. 2019. Pesticide exposure assessment paradigm for solitary bees. *Environmental Entomology* 48: 22–35.
- Sgolastra F, Medrzycki P, Bortolotti L, Maini S, Porrini C, Simon-Delso N, Bosch J. 2020. Bees and pesticide regulation: Lessons from the neonicotinoid experience. *Biodiversity and Conservation* 241: 108356.
- Simon-Delso N, San Martin G, Bruneau E, Hautier L. 2018. Time-to-death approach to reveal chronic and cumulative toxicity of a fungicide for honeybees not revealed with the standard ten-day test. *Scientific Reports* 8: 7241.
- Siviter H, Koricheva J, Brown MJF, Leadbeater E. 2018. Quantifying the impacts of pesticides on learning and memory in bees. *Journal of Applied Ecology* 55: 2812–2821.
- Siviter H, Bailes EJ, Martin CD, Oliver TR, Koricheva J, Leadbetter E, Brown MJF. 2021a. Agrochemicals interact synergistically to increase bee mortality. *Nature* 596: 389–392.
- Siviter H, Richman SK, Muth F. 2021b. Field-realistic neonicotinoid exposure has sub-lethal effects on non-*Apis* bees: A meta-analysis. *Ecology Letters* 24: 2586–2597.
- Smith MR, Singh GM, Mozaffarian D, Myers SS. 2015. Effects of decreases of animal pollinators on human nutrition and global health: A modelling analysis. *Lancet* 386: 1964–1972.

- Sponsler DB, et al. 2019. Pesticides and pollinators: A socioecological synthesis. *Science of the Total Environment* 662: 1012–1027.
- Steffan-Dewenter I, Potts SG, Packer L. 2005. Pollinator diversity and crop pollination services are at risk. *Trends in Ecology and Evolution* 20: 651–652.
- Straw EA, Thompson LJ, Leadbeater E, Brown MJF. 2022. “Inert” ingredients are understudied, potentially dangerous to bees and deserve more research attention. *Proceedings of the Royal Society B* 289: 20212353.
- Tosi S, Nieh JC, Sgolastra F, Cabbri R, Medrzycki P. 2017. Neonicotinoid pesticides and nutritional stress synergistically reduce survival in honey bees. *Proceedings of the Royal Society B* 284: 20171711.
- Tosi S, Nieh JC, Brandt A, Colli M, Fourrier J, Giffard H, Hernández-López J, Malagnini V, Williams GR, Simon-Delso N. 2021. Long-term field-realistic exposure to a next-generation pesticide, flupyradifurone, impairs honey bee behaviour and survival. *Communications Biology* 4: 805.
- Tosi S, Sfeir C, Carnesecchi E, vanEngelsdorp D, Chauzat MP. 2022. Lethal, sublethal, and combined effects of pesticides on bees: A meta-analysis and new risk assessment tools. *Science of the Total Environment* 844: 156857.
- Traynor KS, et al. 2021. Pesticides in honey bee colonies: Establishing a baseline for real world exposure over seven years in the USA. *Environmental Pollution* 279: 116566.
- Trevan JW. 1927. The error of determination of toxicity. *Proceedings of the Royal Society* 101: 483–514.
- Tsvetkov N, Samson-Robert O, Sood K, Patel HS, Malena DA, Gajiwala PH, Maciukiewicz P, Fournier V, Zayed A. 2017. Chronic exposure to neonicotinoids reduces honey bee health near corn crops. *Science* 356: 1395–1397.
- Uhl P, Brühl CA. 2019. The impact of pesticides on flower-visiting insects: A review with regard to European risk assessment. *Environmental Toxicology and Chemistry* 38: 2355–2370.
- [USGS] United States Geological Survey. Pesticide National Synthesis Project. USGS. <https://water.usgs.gov/nawqa/pnsp/usage/maps>.
- Whitehorn PR, O'Connor S, Wackers FL, Goulson D. 2012. Neonicotinoid pesticide reduces bumble bee colony growth and queen production. *Science* 336: 351–352.
- Woodcock BA, Isaac NJB, Bullock JM, Roy DB, Garthwaite DG, Crowe A, Pywell RF. 2016. Impacts of neonicotinoid use on long-term population changes in wild bees in England. *Nature Communications* 7: 12459.
- Woodcock BA, et al. 2017. Country-specific effects of neonicotinoid pesticides on honey bees and wild bees. *Science* 356: 1393–1395.