

ERRATUM

Serotonin Modulates Locomotory Behavior and Coordinates Egg-Laying and Movement in *Caenorhabditis elegans*

Laura Anne Hardaker,¹ Emily Singer,¹ Rex Kerr,¹ Guotong Zhou,² William R. Schafer¹

¹Division of Biology, University of California, San Diego, 9500 Gilman Drive, La Jolla, California 92093-0349

²School of Electrical and Computer Engineering, Georgia Institute of Technology, Atlanta, Georgia 30332-0250

J. Neurobiol. **49**: 303–313, 2001

Table 1 Velocity and Reversal Parameters for Wild-Type, Mutant, and Ablated Animals

Animal Type	Mean Velocity ($\mu\text{m}/\text{sec} \pm \text{S.D.}$)	Mean Reversal Interval ($\text{sec} \pm \text{S.D.}$)	Reversal Parameters	
			μ_1 : Logarithmic Mean Reversal Interval (sec)	σ_1 : Variance (log sec)
N2	15.9 \pm 26.7	37.9 \pm 76.85	17.8	1.19
<i>egl-1</i> (n986)	17.4 \pm 26.9	45.47 \pm 92.24	22.6	1.13
<i>tph-1</i> (mg280)	4.92 \pm 8.4	93.41 \pm 336.45	25.3	1.69
<i>cat-4</i> (e1114)	8.13 \pm 19.3	64.89 \pm 179.16	18.5	1.54
ASH ⁻	16.9 \pm 29.0	40.15 \pm 102.96	19.3	1.17
AVF ⁻	8.54 \pm 15.5	64.01 \pm 131.34	23.6	1.44
<i>glr-1::ICE</i> (AVB ⁻)	8.62 \pm 17.0	172.70 \pm 995.97	19.1	1.86
<i>nmr-1::ICE</i> (AVB ⁺)	7.41 \pm 11.9	198.29 \pm 527.21	42.52	1.75